

Hole Number: ER07-37

Units: METRIC

Project Name: Norway - South Norway	Primary Coordinates Grid: UTM84-32N	Destination Coordinates Grid: UTM:	Collar Dip: -80.90
Project Number: 203	North: 6659511.90	North: 60.07	Collar Az: 58.10
Location: Ertelia	East: 558272.08	East: 10.05	Length: 54.51 (m)
	Elev: 168.89	Elev: 168.89	Start Depth: 0.00 (m)
Date Started: Nov 28, 2007	Collar Survey: N	Plugged: N	Contractor: Drillcon Core AB
Date Completed: Dec 01, 2007	Multishot Survey: N	Hole Size: BQ	Core Storage: Tyrstrand
Logged By: K Leonard	Pulse EM Survey: N	Casing: Left in Hole	Final Depth: 54.51 (m)

Comments: Testing Ni mineralization spatially associated along the Gabbro - Gneiss contact below the Open Cut on Section 1450E. This hole will drill below ER07-36 that intersected a fault zone at 77.6m and was subsequently abandoned.

Results: ER07-37 was abandoned in a FAULT at 54.50m

## Sample Averages

## Survey Data

Depth (m)	Azimuth Decimal	Dip Decimal	Test Type	Flag	Comments	Depth (m)	Azimuth Decimal	Dip Decimal	Test Type	Flag	Comments
10.00	58.10	-80.90	EZ	OK		25.00	59.50	-80.50	EZ	OK	
50.00	59.60	-80.30	EZ	OK							

Detailed Lithology		Assay Data							
From (m)	To (m)	Lithology	Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
0	1.10	CAS, Casing							
1.10	33.40	GNOR, Gabbro Norite dark greenish-grey in colour, medium grained, granular texture, locally sheared, intervening broken core, 1-2% blebby disseminations of Po, sporadic quartz veining, , competent core Mineralization 12.70 - 14.15 : PO Pyrrhotite, BB Blebby, 2% 15.12 - 18.73 : Cpy Chalcopyrite, DIS Disseminated, 0.5% 15.12 - 18.73 : PO Pyrrhotite, BB Blebby, 2% 27.70 - 33.40 : PO Pyrrhotite, BB Blebby, 5% Structure 2.05 - 2.20 limonitic oxide stain 2.49 - 2.75 limonitic oxide stain 4.10 - 5.50 serpentinized, chloritized fractures 19.50 - 20.00 strong biotite halo along both upper and lower contacts. 22.37 - 22.85 : SHR Shear, 45 Deg to CA carbonate, amphibole and biotite altered 29.25 - 29.70 chlorite altered							

Hole Number: ER07-37

Units: METRIC

Detailed Lithology		Lithology	Assay Data						
From (m)	To (m)		Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
33.40	49.87	MGN, Mafic Gneiss Hornblende-Biotite-Chlorite +/- Garnet Gneiss dark grey, medium grained, foliated to sheared, transitional upper contact with GNOR above, cut by two quartz veins with attendant hornblende-biotite alteration around vein contacts, strongly broken core, trace sulphides throughout Structure 35.90 - 37.70 badly broken core, local fault gouge and slickensides 39.30 - 41.90 : VN Veins, 46 Deg to CA milky white vein with 10% included mafic material, locally fractu 47.60 - 48.75 similar to the Qvn observed from 39.30 - 41.90m above, locally badly broken / crushed core							
49.87	54.50	GNOR, Gabbro Norite similar to unit above from 1.10 - 33.40m FLT - strongly fractured, broken core from 53 to 54.50 Structure 53.00 - 54.50 badly broken and crushed core, frequent chlorite slickensides							
54.50	54.51	EOH, End of Hole							