

DETAILED LOG

Hole Number: ER07-39

Units: METRIC

| | | | |
|-------------------------------------|-------------------------------------|------------------------------------|------------------------------|
| Project Name: Norway - South Norway | Primary Coordinates Grid: UTM84-32N | Destination Coordinates Grid: UTM: | Collar Dip: -89.00 |
| Project Number: 203 | North: 6659516.76 | North: 60.07 | Collar Az: 304.00 |
| Location: Ertelia Mine | East: 558195.27 | East: 10.05 | Length: 350.16 (m) |
| | Elev: 167.40 | Elev: 167.40 | Start Depth: 0.00 (m) |
| Date Started: Dec 02, 2007 | Collar Survey: N | Plugged: N | Contractor: Drillcon Core AB |
| Date Completed: Dec 14, 2007 | Multishot Survey: N | Hole Size: BQ | Core Storage: Tyrstrand |
| Logged By: K Leonard | Pulse EM Survey: N | Casing: Left in Hole | Final Depth: 350.16 (m) |

Comments: This hole is designed to test the down dip extension of Ni mineralization intersected in Hole ER07-34. The hole is located about 65m grid south of #34 on Section 1500E.

Results:

Zone 1

88.4 - 89.1m semi-massive @ 50% Po / 0.7m
 89.1 - 90.32m disseminated @ 3-5% / 1.22m

Zone 2

114.38 - 118.90m semi-massive @ 65% Po / 4.52m
 118.90 - 120.70m late rubbly FLT Zone @ <5% Po
 120.70 - 122.10m disseminated Po @ 10-15%, minor Cpy
 122.10 - 125.00m semi-massive @ 50% Po, Cpy/ 2.9m (net-textured, frags, Bx)

Sample Averages

| Average Type | From (m) | To (m) | Length (m) | Ni% | Cu% | Co% |
|--------------|----------|--------|------------|--------|--------|--------|
| WEIGHTED | 88.35 | 89.07 | 0.72 | 0.8593 | 0.3632 | 0.0764 |
| WEIGHTED | 114.38 | 118.90 | 4.52 | 0.6362 | 0.7526 | 0.0461 |
| WEIGHTED | 114.38 | 125.00 | 10.62 | 0.4942 | 0.6610 | 0.0419 |
| WEIGHTED | 120.70 | 125.00 | 4.30 | 0.5375 | 0.7965 | 0.0528 |

Survey Data

| Depth (m) | Azimuth Decimal | Dip Decimal | Test Type | Flag | Comments | Depth (m) | Azimuth Decimal | Dip Decimal | Test Type | Flag | Comments |
|-----------|--------------------|----------------|--------------|------|----------|-----------|--------------------|----------------|--------------|------|----------|
| 10.00 | | -89.00 | EZ | OK | | 25.00 | 306.50 | -89.50 | EZ | OK | |
| 50.00 | 327.70 | -89.60 | EZ | OK | | 100.00 | 302.00 | -89.30 | EZ | OK | |

| Detailed Lithology | | Assay Data | | | | | | | |
|--------------------|--------|-------------|---------------|----------|--------|------------|-----|-----|-----|
| From (m) | To (m) | Lithology | Sample Number | From (m) | To (m) | Length (m) | Ni% | Cu% | Co% |
| 0 | 4.90 | CAS, Casing | | | | | | | |

Hole Number: ER07-39

Units: METRIC

| Detailed Lithology | | Lithology | Assay Data | | | | | | |
|--------------------|--------|--|---------------|----------|--------|------------|-----|-----|-----|
| From (m) | To (m) | | Sample Number | From (m) | To (m) | Length (m) | Ni% | Cu% | Co% |
| 4.90 | 30.00 | <p>GNOR, Gabbro Norite grey in colour, medium grained, massive to weakly foliated to locally sheared, cut by narrow mafic dykes,</p> <p>composed of plagioclase, orthopyroxene, hornblende / actinolite? and accessory quartz, chlorite, carbonate, garnet, olivine? and serpentinite, variably biotite and amphibole-rich where sheared, intermittent blebby Po, competent core., faulted lower contact with PEG.</p> <p>Structure 6.40 - 6.90 : F Fractured, 20 Deg to CA 10.65 - 10.90 : F Fractured, 10 Deg to CA 17.25 - 18.00 : F Fractured, 5 Deg to CA slickensides, cherty, apple green alteration 19.85 - 20.00 apple green carbonate 20.28 - 20.48 apple green carbonate 29.90 - 30.00 gouge, brecciation and shearing</p> <p>MINOR INTERVALS: Minor Interval: 13.2 - 13.37 MD, Mafic Dike grey, fine grained, sharp upper and lower contacts at 70 deg and 65 deg to the LCA respectively Minor Interval: 13.9 - 14.7 MD, Mafic Dike same as unit observed from 13.20 to 13.37m, nil sulphides</p> <p>upper contact trends 40 deg to the LCA. lower contact oriented at 52 deg to the LCA. Minor Interval: 15.95 - 16.2 MD, Mafic Dike same as intersected above, broken upper contact and sharp lower contact at 43 deg to the LCA.</p> | | | | | | | |
| 30.00 | 37.05 | <p>PEG, Pegmatite mottled black and white, predominately silicified with 20-30% included mafic material, sheared at the upper contact with GNOR at 70 deg to the LCA, becomes more mafic-rich down section, locally brecciated, nil sulphides</p> <p>the unit is interrupted by inclusions of coarse grained garnetiferous mafic gneiss from 33.85 - 34.50m, 35.32 - 35.92m and 36.25 - 36.50m.</p> | | | | | | | |
| 37.05 | 38.70 | <p>MGN, Mafic Gneiss dark grey, coarse grained, foliated to sheared, abundant hornblende, garnet and accessory biotite, altered GABB / GNOR immediately below the lower contact with PEG unit, moderately broken / blocky core, nil sulphides</p> | | | | | | | |

Hole Number: ER07-39

Units: METRIC

| Detailed Lithology | | Lithology | Assay Data | | | | | | |
|--------------------|--------|--|---------------|----------|--------|------------|--------|--------|--------|
| From (m) | To (m) | | Sample Number | From (m) | To (m) | Length (m) | Ni% | Cu% | Co% |
| 38.70 | 82.57 | GNOR, Gabbro Norite similar to unit intersected above from 4.90 - 30.00m Structure 49.70 - 49.90 51.20 - 52.90 slickensides, vitreous apple green carbonate alteration 58.00 - 78.65 : SHR Shear, 50 Deg to CA narrow shear band, chlorite-carbonate altered 66.00 - 66.90 : F Fractured, 10 Deg to CA striated, slickensides, moderate carbonate alteration 81.58 - 81.65 : SHR Shear, 50 Deg to CA chlorite-carbonate altered 82.53 - 82.57 : SHR Shear, 34 Deg to CA chlorite-carbonate altered MINOR INTERVALS: Minor Interval: 80.95 - 81.1 PEG, Pegmatite narrow seam of milky white quartz with 20% included mafic material as inclusions and aligned along healed fractures, nil sulphides. 8-10cm wide corona above upper contact consisting of garnet and comminuted amphibole and biotite. sharp upper contact at 45 deg to the LCA. sharp lower contact at 65 deg to the LCA | QB3674 | 81.50 | 82.00 | 0.50 | 0.0300 | 0.0070 | 0.0040 |
| | | | QB3675 | 82.00 | 82.50 | 0.50 | 0.0330 | 0.0170 | 0.0040 |
| | | | QB3676 | 82.50 | 83.00 | 0.50 | 0.1850 | 0.1380 | 0.0160 |
| 82.57 | 84.60 | SULF, Sulfide tr to 5-7% disseminated, blebby Po, Py and minor Cpy form irregular coarse clots, dense f.g. masses and discrete grains, some internal fracturing of grains | QB3677 | 83.00 | 83.50 | 0.50 | 0.1140 | 0.0980 | 0.0100 |
| | | | QB3678 | 83.50 | 83.90 | 0.40 | 0.1790 | 0.1520 | 0.0160 |
| | | | QB3679 | 83.90 | 84.30 | 0.40 | 0.1110 | 0.0220 | 0.0060 |
| | | | QB3681 | 84.30 | 84.65 | 0.35 | 0.0700 | 0.0100 | 0.0040 |
| 84.60 | 88.07 | GNOR, Gabbro Norite similar to units above from 4.90 - 30.0m and 82.57 - 84.60m, sparse, fine to-medium grained sulphides. | QB3682 | 84.65 | 84.90 | 0.25 | 0.0770 | 0.0650 | 0.0080 |
| | | | PG05218 | 84.90 | 85.25 | 0.35 | 0.0790 | 0.0550 | 0.0070 |
| | | | QB3683 | 85.25 | 85.60 | 0.35 | 0.0450 | 0.0160 | 0.0040 |
| | | | QB3684 | 85.60 | 86.10 | 0.50 | 0.0290 | 0.0080 | 0.0020 |
| | | | QB3685 | 86.10 | 87.00 | 0.90 | 0.0290 | 0.0080 | 0.0020 |
| | | | QB3686 | 87.00 | 87.50 | 0.50 | 0.0290 | 0.0070 | 0.0020 |
| | | | QB3687 | 87.50 | 88.00 | 0.50 | 0.0510 | 0.0120 | 0.0030 |
| | | | QB3688 | 88.00 | 88.35 | 0.35 | 0.1160 | 0.0640 | 0.0100 |
| 88.07 | 90.32 | SULF, Sulfide disseminated to semi-massive sulphides - Po >Py>Cpy centered at 88.50m 88.31-89.05m +50% semi-massive Po>Py and minor Cpy 89.05-90.32m 3-5% disseminated, blebby Po | QB3689 | 88.35 | 88.70 | 0.35 | 1.1800 | 0.2090 | 0.1030 |
| | | | QB3690 | 88.70 | 89.07 | 0.37 | 0.5560 | 0.5090 | 0.0510 |
| | | | QB3691 | 89.07 | 89.50 | 0.43 | 0.1060 | 0.0690 | 0.0090 |
| | | | QB3692 | 89.50 | 90.00 | 0.50 | 0.0800 | 0.0160 | 0.0040 |
| | | | QB3693 | 90.00 | 90.50 | 0.50 | 0.0570 | 0.0180 | 0.0030 |

Hole Number: ER07-39

Units: METRIC

| Detailed Lithology | | Lithology | Assay Data | | | | | | |
|--------------------|--------|--|---------------|----------|--------|------------|--------|--------|--------|
| From (m) | To (m) | | Sample Number | From (m) | To (m) | Length (m) | Ni% | Cu% | Co% |
| 90.32 | 114.38 | GNOR, Gabbro Norite grey in colour, medium to- coarse grained, massive to locally sheared, intermittent blebby Po from 97.0 - 97.50m - 5 - 10% coarse grained, disseminated Po > Py and minor Cpy "chock a block" with amphibole and pyroxene crystals - hypidiomorphic texture. Structure 102.50 - 105.55 fractured, blocky core from < 2 to 20cm segments. MINOR INTERVALS: Minor Interval: 101.89 - 102.05 MD, Mafic Dike narrow seam with indistinct contacts black in colour, aphanitic texture, nil sulphides | QB3694 | 90.50 | 91.00 | 0.50 | 0.0430 | 0.0090 | 0.0030 |
| | | | QB3695 | 113.90 | 114.38 | 0.48 | 0.0250 | 0.0160 | 0.0020 |
| 114.38 | 118.90 | SMS, Semi Massive Sulphide 60-65% semi-massive to massive Po < Py < Cpy mineralization, semi-massive mineralization is net-textured and brecciated, conspicuous dark grey coloured wallrock fragments | QB3696 | 114.38 | 114.85 | 0.47 | 0.9610 | 0.1910 | 0.0740 |
| | | | QB3697 | 114.85 | 115.15 | 0.30 | 1.3530 | 0.1470 | 0.0980 |
| | | | QB3698 | 115.15 | 115.55 | 0.40 | 0.2960 | 0.5260 | 0.0200 |
| | | | QB3699 | 115.55 | 116.00 | 0.45 | 0.2800 | 0.9610 | 0.0220 |
| | | | QB3701 | 116.00 | 116.50 | 0.50 | 0.3880 | 1.2510 | 0.0280 |
| | | | QB3702 | 116.50 | 117.00 | 0.50 | 0.6890 | 1.4410 | 0.0460 |
| | | | QB3703 | 117.00 | 117.35 | 0.35 | 0.5910 | 0.7320 | 0.0440 |
| | | | QB3704 | 117.35 | 117.85 | 0.50 | 0.6930 | 0.7540 | 0.0480 |
| | | | QB3705 | 117.85 | 118.30 | 0.45 | 1.1350 | 0.1390 | 0.0830 |
| | | | QB3706 | 118.30 | 118.90 | 0.60 | 0.2850 | 0.9720 | 0.0210 |
| 118.90 | 120.70 | FLT, Fault crushed, blocky core - GNOR, blocky pieces range in size from <1 to 20cm in length. between trace and 5% disseminated sulphides | QB3707 | 118.90 | 119.60 | 0.70 | 0.0330 | 0.1740 | 0.0050 |
| | | | QB3708 | 119.60 | 120.00 | 0.40 | 0.0480 | 0.0290 | 0.0060 |
| | | | QB3709 | 120.00 | 120.70 | 0.70 | 0.0280 | 0.0860 | 0.0050 |
| 120.70 | 122.00 | SULF, Sulfide 10 to 15% disseminated and net-textured Po > Py and minor Cpy 121.08 - 121.60m - bleached core, minor quartz veins, 25% net-textured sulphides | QB3710 | 120.70 | 121.10 | 0.40 | 0.4630 | 0.2440 | 0.0350 |
| | | | QB3711 | 121.10 | 121.60 | 0.50 | 0.3520 | 0.1160 | 0.0220 |
| | | | QB3712 | 121.60 | 122.00 | 0.40 | 0.0820 | 0.0860 | 0.0060 |
| 122.00 | 125.00 | SMS, Semi Massive Sulphide net-textured and brecciated + 50% semi-massive Po > Py > Cpy. From millimeter to 25cm fragments throughout the mineralization. Cpy ranges locally from 5 to 8% from 122.75 - 122.90; 124.0 - 124.17m and 124.75 - 124.80m. | QB3713 | 122.00 | 122.50 | 0.50 | 0.3080 | 0.8550 | 0.0760 |
| | | | QB3714 | 122.50 | 123.00 | 0.50 | 0.5910 | 1.1730 | 0.0940 |
| | | | QB3715 | 123.00 | 123.45 | 0.45 | 1.1430 | 0.3830 | 0.0760 |
| | | | QB3716 | 123.45 | 123.90 | 0.45 | 0.0320 | 0.0340 | 0.0030 |
| | | | QB3717 | 123.90 | 124.40 | 0.50 | 0.9680 | 3.4650 | 0.0600 |
| | | | QB3718 | 124.40 | 125.00 | 0.60 | 0.7580 | 0.5010 | 0.0820 |

Hole Number: ER07-39

Units: METRIC

| Detailed Lithology | | Lithology | Assay Data | | | | | | |
|--------------------|--------|--|---------------|----------|--------|------------|--------|--------|--------|
| From (m) | To (m) | | Sample Number | From (m) | To (m) | Length (m) | Ni% | Cu% | Co% |
| 125.00 | 291.65 | FGN, Felsic Gneiss FOOTWALL FGN grey to variably coloured, fine to medium grained, gneissic texture, conspicuously laminated / banded at 35 deg to the LCA, colour laminated, composed of granular quartz-feldspar-biotitite+/-hornblende+/-garnet+/-chlorite schist. 254.0 - 264.50m - garnetiferous section, isolated subhedra (0.5mm - 3cm) and f.g masses and bands. indistinct, gradational contact with well mineralized GNOR unit above! Mineralization 127.90 - 128.11 : Cpy Chalcopyrite, F Fracture Controlled, 3% irregular splashes 127.90 - 128.11 : PO Pyrrhotite, F Fracture Controlled, 30% 30% remobilized Po-Cpy associated with smokey grey quartz veinlet Structure 129.00 - 130.05 151.10 - 152.50 : F Fractured, 45 Deg to CA 157.60 - 157.80 : LAM Laminated, 35 Deg to CA 165.00 - 165.10 : LAM Laminated, 33 Deg to CA 172.10 - 173.00 strongly broken core 174.30 - 174.55 strongly broken core 176.00 - 176.55 : LAM Laminated, 34 Deg to CA 180.00 - 180.30 well fractured interval 201.20 - 201.33 : LAM Laminated, 34 Deg to CA 204.80 - 206.10 : F Fractured, 10 Deg to CA chlorite slickensides 208.20 - 208.60 : F Fractured, 5 Deg to CA chlorite altered fracture faces 218.25 - 218.90 236.40 - 241.00 246.75 - 246.81 : VN Veins, 40 Deg to CA small scale dextral offset of quartz stringers 262.90 - 263.00 : FOL Foliated, 37 Deg to CA 271.90 - 274.50 : GN Gneissic, 30 Deg to CA prominent gneissic banding - colour layering 276.74 - 278.66 : GN Gneissic, 33 Deg to CA 287.30 - 290.92 : GN Gneissic, 40 Deg to CA strongly fractured section | QB3719 | 125.00 | 125.50 | 0.50 | 0.0760 | 0.0210 | 0.0070 |
| | | | QB3721 | 125.50 | 127.00 | 1.50 | 0.0720 | 0.0720 | 0.0070 |
| | | | QB3727 | 127.00 | 127.40 | 0.40 | 0.0480 | 0.0440 | 0.0040 |
| | | | QB3722 | 127.40 | 128.10 | 0.70 | 0.5460 | 0.3330 | 0.0440 |
| | | | QB3723 | 128.10 | 128.50 | 0.40 | 0.0400 | 0.0390 | 0.0040 |

DETAILED LOG

Hole Number: ER07-39

Units: METRIC

| Detailed Lithology | | Assay Data | | | | | | | |
|--------------------|--------|---|---------------|----------|--------|------------|-----|-----|-----|
| From (m) | To (m) | Lithology | Sample Number | From (m) | To (m) | Length (m) | Ni% | Cu% | Co% |
| | | <p>MINOR INTERVALS: Minor Interval: 125 - 127 MD, Mafic Dike similar to mafic dyke units observed above</p> <p>dark grey, fine grained, homogenous, indistinct upper and lower contacts, nil sulphides</p> <p>Minor Interval: 143 - 148 PEG, Pegmatite intercalated PEG and FGN</p> <p>PEG is predominately mottled quartz and 25-40% included mafic material.</p> | | | | | | | |
| 291.65 | 307.00 | <p>IGN, Intermediate Gneiss grey in colour, medium grained, layered gneissic texture (colour banding), consists of quartz-plagioclase-hornblende-biotite- garnet schist, conspicuously more amphibole than that observed above from 125 - 291.65m.</p> <p>Structure 291.65 - 307.00 : GN Gneissic, 31 Deg to CA distinct colour banding</p> | | | | | | | |

Hole Number: ER07-39

Units: METRIC

| Detailed Lithology | | Lithology | Assay Data | | | | | | |
|--------------------|--------|--|---------------|----------|--------|------------|-----|-----|-----|
| From (m) | To (m) | | Sample Number | From (m) | To (m) | Length (m) | Ni% | Cu% | Co% |
| 307.00 | 350.15 | FGN, Felsic Gneiss similar to unit observed from 125 to 291.65m above, strongly tectonized unit - widespread brecciation, nil sulphides, competent core Structure 311.00 - 313.00 cm-scale quartz fragments 313.00 - 315.45 strong brecciation, chlorite alteration, fissile fault gouge 319.00 - 319.45 321.76 - 323.32 strongly bleached, cherty core, stockworked quartz veinlets 324.00 - 327.38 : FOL Foliated, 44 Deg to CA gneissic layering 332.45 - 334.25 intensely fractured 336.74 - 338.81 stockworked, ptygmatic veinlets 338.45 - 339.75 irregular carbonate stringers 338.45 - 339.75 fractured, graphitic 339.85 - 339.95 : SHR Shear, 38 Deg to CA soft, black and graphitic 350.00 - 350.15 : GN Gneissic, 32 Deg to CA colour laminated | | | | | | | |
| 350.15 | 350.16 | EOH, End of Hole | | | | | | | |

Samples

| Sample Number | From (m) | To (m) | Ni% | Cu% | Co% |
|---------------|----------|--------|--------|--------|--------|
| Sample Type | ASSAY | | | | |
| QB3674 | 81.50 | 82.00 | 0.0300 | 0.0070 | 0.0040 |
| QB3675 | 82.00 | 82.50 | 0.0330 | 0.0170 | 0.0040 |
| QB3676 | 82.50 | 83.00 | 0.1850 | 0.1380 | 0.0160 |
| QB3677 | 83.00 | 83.50 | 0.1140 | 0.0980 | 0.0100 |
| QB3678 | 83.50 | 83.90 | 0.1790 | 0.1520 | 0.0160 |
| QB3679 | 83.90 | 84.30 | 0.1110 | 0.0220 | 0.0060 |
| QB3681 | 84.30 | 84.65 | 0.0700 | 0.0100 | 0.0040 |
| QB3682 | 84.65 | 84.90 | 0.0770 | 0.0650 | 0.0080 |
| PG05218 | 84.90 | 85.25 | 0.0790 | 0.0550 | 0.0070 |
| QB3683 | 85.25 | 85.60 | 0.0450 | 0.0160 | 0.0040 |
| QB3684 | 85.60 | 86.10 | 0.0290 | 0.0080 | 0.0020 |
| QB3685 | 86.10 | 87.00 | 0.0290 | 0.0080 | 0.0020 |

Hole Number: ER07-39

Units: METRIC

Samples

| Sample Number | From (m) | To (m) | Ni% | Cu% | Co% |
|---------------|----------|--------|--------|--------|--------|
| Sample Type | ASSAY | | | | |
| QB3686 | 87.00 | 87.50 | 0.0290 | 0.0070 | 0.0020 |
| QB3687 | 87.50 | 88.00 | 0.0510 | 0.0120 | 0.0030 |
| QB3688 | 88.00 | 88.35 | 0.1160 | 0.0640 | 0.0100 |
| QB3689 | 88.35 | 88.70 | 1.1800 | 0.2090 | 0.1030 |
| QB3690 | 88.70 | 89.07 | 0.5560 | 0.5090 | 0.0510 |
| QB3691 | 89.07 | 89.50 | 0.1060 | 0.0690 | 0.0090 |
| QB3692 | 89.50 | 90.00 | 0.0800 | 0.0160 | 0.0040 |
| QB3693 | 90.00 | 90.50 | 0.0570 | 0.0180 | 0.0030 |
| QB3694 | 90.50 | 91.00 | 0.0430 | 0.0090 | 0.0030 |
| QB3695 | 113.90 | 114.38 | 0.0250 | 0.0160 | 0.0020 |
| QB3696 | 114.38 | 114.85 | 0.9610 | 0.1910 | 0.0740 |
| QB3697 | 114.85 | 115.15 | 1.3530 | 0.1470 | 0.0980 |
| QB3698 | 115.15 | 115.55 | 0.2960 | 0.5260 | 0.0200 |
| QB3699 | 115.55 | 116.00 | 0.2800 | 0.9610 | 0.0220 |
| QB3701 | 116.00 | 116.50 | 0.3880 | 1.2510 | 0.0280 |
| QB3702 | 116.50 | 117.00 | 0.6890 | 1.4410 | 0.0460 |
| QB3703 | 117.00 | 117.35 | 0.5910 | 0.7320 | 0.0440 |
| QB3704 | 117.35 | 117.85 | 0.6930 | 0.7540 | 0.0480 |
| QB3705 | 117.85 | 118.30 | 1.1350 | 0.1390 | 0.0830 |
| QB3706 | 118.30 | 118.90 | 0.2850 | 0.9720 | 0.0210 |
| QB3707 | 118.90 | 119.60 | 0.0330 | 0.1740 | 0.0050 |
| QB3708 | 119.60 | 120.00 | 0.0480 | 0.0290 | 0.0060 |
| QB3709 | 120.00 | 120.70 | 0.0280 | 0.0860 | 0.0050 |
| QB3710 | 120.70 | 121.10 | 0.4630 | 0.2440 | 0.0350 |
| QB3711 | 121.10 | 121.60 | 0.3520 | 0.1160 | 0.0220 |
| QB3712 | 121.60 | 122.00 | 0.0820 | 0.0860 | 0.0060 |
| QB3713 | 122.00 | 122.50 | 0.3080 | 0.8550 | 0.0760 |
| QB3714 | 122.50 | 123.00 | 0.5910 | 1.1730 | 0.0940 |
| QB3715 | 123.00 | 123.45 | 1.1430 | 0.3830 | 0.0760 |
| QB3716 | 123.45 | 123.90 | 0.0320 | 0.0340 | 0.0030 |
| QB3717 | 123.90 | 124.40 | 0.9680 | 3.4650 | 0.0600 |
| QB3718 | 124.40 | 125.00 | 0.7580 | 0.5010 | 0.0820 |
| QB3719 | 125.00 | 125.50 | 0.0760 | 0.0210 | 0.0070 |
| QB3721 | 125.50 | 127.00 | 0.0720 | 0.0720 | 0.0070 |
| QB3727 | 127.00 | 127.40 | 0.0480 | 0.0440 | 0.0040 |
| QB3722 | 127.40 | 128.10 | 0.5460 | 0.3330 | 0.0440 |

Hole Number: ER07-39

Units: METRIC

Samples

| Sample Number | From (m) | To (m) | Ni% | Cu% | Co% |
|-------------------|----------|--------|--------|--------|--------|
| Sample Type ASSAY | | | | | |
| QB3723 | 128.10 | 128.50 | 0.0400 | 0.0390 | 0.0040 |