

DETAILED LOG

Hole Number: ES07-56

Units: Metric

Project Name: Norway - Espedalen	Primary Coordinates Grid: UTM84-32N	Destination Coordinates Grid: UTM:	Collar Dip: -51.50
Project Number: 201	North: 6801309.79	North: 61.34	Collar Az: 236.50
Location: Stormyra	East: 535140.50	East: 9.66	Length: 152.00 (m)
	Elev: 975.34	Elev: 975.34	Start Depth: 0.00 (m)
Date Started: Mar 26, 2007	Collar Survey: Y	Plugged: N	Contractor: Geo Drilling A/S
Date Completed: Mar 28, 2007	Multishot Survey: N	Hole Size: TT46	Core Storage:
Logged By: jdnor	Pulse EM Survey: N	Casing: Left in hole. capped.	Final Depth: 152.00 (m)

Comments: Hole designed to test 25m down dip of mineralized zone intersected in ES2004-08. Will test botom edge of UTEM plate (surface survey).

Sample Averages

Average Type	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
WEIGHTED	85.50	87.00	1.50	0.6333	0.2321	0.0258

Detailed Lithology			Assay Data						
From (m)	To (m)	Lithology	Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
0	5.80	CAS, Casing							

Hole Number: ES07-56

Units: Metric

Detailed Lithology		Lithology	Assay Data						
From (m)	To (m)		Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
5.80	45.60	<p>ANOR, Anorthosite</p> <p>ANORTHOSITE</p> <p>Very light green to white and fine to medium grained.</p> <p>Pale green sericitized feldspar?</p> <p>White to cream coloured phases less altered feldspar and quartz.</p> <p>Mixed medium green-grey mafic dykes? - generally thin (<0.20m) and often contain anorthositic wall-rock fragments.</p> <p>Dark green chloritic bands and blebs, appear to be aligned parallel to main fabric.</p> <p>Foliation or main fabric initially at 85 deg. to CA, but can flatten locally to 50 deg.</p> <p>Unit is not mineralized.</p> <p>Alteration</p> <p>5.80 - 44.60 :Ser Sericite, P Pervasive, M Moderate</p> <p>5.80 - 44.60 :CHL Chlorite, H Patchy, W Weak</p> <p>Structure</p> <p>6.60 - 6.62 : FOL Foliated, 80 Deg to CA marked by chloritic band</p> <p>13.10 - 13.12 : FOL Foliated, 70 Deg to CA</p> <p>18.30 - 18.32 : FOL Foliated, 65 Deg to CA</p> <p>35.70 - 35.72 : FOL Foliated, 70 Deg to CA</p> <p>42.10 - 42.12 : FOL Foliated, 75 Deg to CA</p> <p>MINOR INTERVALS:</p> <p>Minor Interval:</p> <p>9.18 - 9.48 MD, Mafic Dike</p> <p>Mafic Dyke</p> <p>UC undulating at 65 to 85 deg. to CA.</p> <p>Medium green-grey, fine grained and granular.</p> <p>5% mixed anorthositic wall-rock fragments.</p> <p>LC irregular - sheared?</p>							
45.60	52.60	<p>MD, Mafic Dike</p> <p>MAFIC DYKE</p> <p>Upper contact sharp at 60 to 65 deg. to CA</p> <p>Medium green, fine grained and relatively massive.</p> <p>Weak fabric at 70 to 80 deg. - marked by lighter coloured felspar rich bands/zones.</p> <p>Unit is not mineralized.</p> <p>Lower contact at 70 to 80 deg. to CA.</p> <p>Structure</p> <p>45.60 - 45.61 : UC Upper Contact, 60 Deg to CA</p> <p>52.59 - 52.60 : LC Lower Contact, 75 Deg to CA</p>							

Hole Number: ES07-56

Units: Metric

Detailed Lithology		Lithology	Assay Data						
From (m)	To (m)		Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
52.60	61.40	<p>ANOR, Anorthosite</p> <p>ANORTHOSITE</p> <p>Similar to initial unit.</p> <p>Predominantly a pale green with white to cream coloured streaks and bands.</p> <p>Moderately to strongly sericitic throughout.</p> <p>Local, but minor dark green chloritic wisps and blebs.</p> <p>Local hematite staining - from pink to dark red on foliation surfaces.</p> <p>Fabric or foliation varies, locally compositonal banding hoighly deformed, but at approximately 80 to 85 deg. to CA.</p> <p>59.90 - 61.40 Weak patchy hematitic staining Fabric (foliation) varies from 50 to 70 deg., and is often contorted (folded?).</p> <p>Alteration</p> <p>59.90 - 61.40 :HE Hematite, PCH Patchy, W Weak</p> <p>52.60 - 61.40 :Ser Sericite, P Pervasive, M Moderate</p> <p>52.60 - 61.40 :CH Chlorite, H Patchy, W Weak</p> <p>Structure</p> <p>53.70 - 53.71 : FOL Foliated, 80 Deg to CA</p> <p>57.40 - 57.41 : FOL Foliated, 65 Deg to CA marked by chloritic banding.</p> <p>61.20 - 61.20 : FOL Foliated, 50 Deg to CA sheared?</p>							
61.40	62.55	<p>MD, Mafic Dike</p> <p>MAFIC DYKE</p> <p>Upper contact sheared, but at 60 deg. to CA.</p> <p>Medium to dark grey-green and weakly foliated at 75 deg. to CA.</p> <p>Local inclusions anorthosite.</p> <p>Unit is not mineralized.</p> <p>Lower contact undulating at 35 to 50 deg. to CA.</p> <p>Structure</p> <p>61.40 - 61.41 : UC Upper Contact, 60 Deg to CA</p> <p>62.54 - 62.55 : LC Lower Contact, 45 Deg to CA</p>							

Hole Number: ES07-56

Units: Metric

Detailed Lithology		Lithology	Assay Data						
From (m)	To (m)		Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
62.55	73.10	ANOR, Anorthosite	PG07001	62.55	63.20	0.65	0.0050	0.0025	0.0030
		ANORTHOSITE	PG07002	63.20	63.50	0.30	0.0070	0.0060	0.0050
		Unit mixed with several sheared mafic and ultramafic dykes, the initial two of which are mineralized.	PG07003	63.50	64.62	1.12	0.0040	0.0025	0.0030
		The Anorthosite varies from pale green to cream coloured and locally redish due to hematite staining.	PG07004	64.62	64.98	0.36	1.2250	0.4170	0.0350
		Strongly sheared/tectonized throughout.	PG07005	64.98	65.70	0.72	0.0100	0.0025	0.0030
		Anorthosite moderately to strongly sericitized throughout.							
		Minor chloitic bands/wisps.							
		62.55 - 63.29 Weak to moderate patchy hematitic alteration - interstial.							
		63.50 - 64.62 Weak patchy hematite alteration.							
		64.98 - 66.00 Moderate interstitial hematite alteration. Fabric varies, but predominantly at 80 to 85 deg. to CA.							
		Alteration							
		62.55 - 73.10 :Ser Sericite, P Pervasive, M Moderate							
		63.50 - 64.62 :HE Hematite, H Patchy, W Weak							
		64.98 - 66.00 :HE Hematite, INT Interstitial, M Moderate							
		62.55 - 63.29 :HE Hematite, H Patchy, W Weak							
		MINOR INTERVALS:							
		Minor Interval:							
		63.29 - 63.5 UM, Ultramafic							
		Sheared Ultramafic							
		Upper contact at 45 deg. to CA.							
		Medium to dark green and fine grained.							
		Dark green phase serpentinous.							
		2% to 3% fine fracture controlled Py.							
		Lower contact at 40 to 45 deg. to CA.							
		Mineralization							
		63.29 - 63.50 : PY Pyrite, DIS Disseminated, 2%							
		Alteration							
		63.29 - 63.50 :SRP Serpentine, P Pervasive, W Weak							
		Structure							
		63.29 - 63.30 : UC Upper Contact, 45 Deg to CA							
		63.49 - 63.50 : LC Lower Contact, 40 Deg to CA							

Hole Number: ES07-56

Units: Metric

Detailed Lithology		Assay Data							
From (m)	To (m)	Lithology	Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
		<p>MINOR INTERVALS:</p> <p>Minor Interval:</p> <p>64.62 - 64.98 MD, Mafic Dike</p> <p>Sheared Mafic Dyke</p> <p>Upper contact badly broken.</p> <p>Medium to dark green, fine grained and badly broken.</p> <p>Overall carries 15% sulphides as thin stringers and fracture fillings?</p> <p>Most sulphides concentrated about lower contact - 5 to 15mm semi-massive stringer.</p> <p>Sulphides: 10% Po, 4% Py and 1% Cpy.</p> <p>Lower contact at 35 to 40 deg. to CA.</p> <p>Mineralization</p> <p>64.62 - 64.98 : PO Pyrrhotite, STR Stringers, 10%</p> <p>64.62 - 64.98 : Cpy Chalcopyrite, STR Stringers, 1%</p> <p>64.62 - 64.98 : PY Pyrite, STR Stringers, 4%</p> <p>Structure</p> <p>64.62 - 64.63 : LC Lower Contact, 40 Deg to CA</p> <p>Minor Interval:</p> <p>68.2 - 68.7 MD, Mafic Dike</p> <p>Mafic Dyke</p> <p>Upper Contact at 75 to 80 deg. to CA.</p> <p>Medium to dark green, fine grained and weakly tectonized/sheared.</p> <p>Minor fracture controlled Py adjacent to lower contact.</p> <p>Lower contact at 80 deg. to CA.</p> <p>Structure</p> <p>68.20 - 68.21 : UC Upper Contact, 75 Deg to CA</p> <p>68.69 - 68.70 : LC Lower Contact, 80 Deg to CA</p> <p>Minor Interval:</p> <p>71.36 - 72.12 UM, Ultramafic</p> <p>Sheared Ultramafic</p> <p>Upper contact at approx. 80 deg. to CA.</p> <p>Dark grey-green, fine grained and soft.</p> <p>Moderately serpentinous.</p> <p>20% anorthositic wall-rock inclusions.</p> <p>Unit is not mineralized.</p> <p>Lowercontact at 60 deg. to CA.</p> <p>Alteration</p> <p>71.36 - 72.12 :SERP Serpentine, P Pervasive, M Moderate</p> <p>Structure</p> <p>71.36 - 71.37 : LC Lower Contact, 60 Deg to CA</p> <p>71.36 - 71.37 : UC Upper Contact, 80 Deg to CA</p>							

DETAILED LOG

Hole Number: ES07-56

Units: Metric

Detailed Lithology		Assay Data							
From (m)	To (m)	Lithology	Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
73.10	75.90	MD, Mafic Dike MAFIC DYKE Upper contact sharp at 70 deg. to CA. Medium green, fine grained with a weak fabric at 70 to 80 deg. to CA. Fine white speckled feldspars? throughout - often forming crude bands paralleling foliation. Unit is not mineralized. Lower contact sharp at 80 deg. to CA. Structure 73.10 - 73.11 : LC Lower Contact, 80 Deg to CA 73.10 - 73.11 : UC Upper Contact, 70 Deg to CA							

Hole Number: ES07-56

Units: Metric

Detailed Lithology		Assay Data							
From (m)	To (m)	Lithology	Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
75.90	99.60	ANOR, Anorthosite ANORTHOSITE Strongly deformed and altered. Colour varies from pale green/white to dark red/white. Moderately to strongly sericitic throughout. Patches and bands chlorite - patchy. Local hematite stained sections - red to pinkish. Well foliated - often contorted, but main orientation at 80 to 85 deg. to CA.	PG07006	84.00	85.00	1.00	0.0280	0.0330	0.0030
			PG07007	85.00	85.50	0.50	0.0120	0.0150	0.0030
			PG07008	85.50	85.95	0.45	0.7140	0.2390	0.0260
			PG07009	85.95	86.40	0.45	0.1590	0.1490	0.0100
			PG07010	86.40	86.70	0.30	1.6400	0.3000	0.0660
			PG07012	86.70	87.00	0.30	0.2170	0.2780	0.0090
			PG07013	87.00	88.00	1.00	0.0720	0.1200	0.0050
		77.80 - 80.45 Mixed red/green section - moderately to strongly hematite stained section. Very distinct. Well foliated at 75 to 80 deg. to CA.							
		84.00 - 85.53 Section with 2% to 3% fine dark speckling - dark brown under hand lens - biotite? Minor fine disseminated Cpy.							
		95.04 - 97.20 Weakly hematitic stained section. Local fractures strongly hematitic.							
		Alteration 95.04 - 97.20 :HE Hematite, H Patchy, W Weak 77.80 - 80.45 :HE Hematite, P Pervasive, M Moderate 84.00 - 85.53 :BIO Biotite, SP Spotted, W Weak 75.90 - 99.60 :Ser Sericite, P Pervasive, M Moderate							
		Structure 83.70 - 83.71 : FOL Foliated, 70 Deg to CA							
		MINOR INTERVALS: Minor Interval: 76.12 - 76.4 MD, Mafic Dike Mafic Dyke Uper contact at 75 deg. to CA - not sharp. Medium to dark green and fine grained. Partial UM component? Well foliated - locally finely contorted, at 70 to 75 deg. to CA. Unit is not mineralized. Lower contact sharp at 70 deg - sheared?							
		Structure 76.12 - 76.13 : UC Upper Contact, 75 Deg to CA 76.39 - 76.40 : LC Lower Contact, 70 Deg to CA							

Hole Number: ES07-56

Units: Metric

Detailed Lithology		Assay Data							
From (m)	To (m)	Lithology	Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
		MINOR INTERVALS: Minor Interval: 85.53 - 87.08 MD, Mafic Dike Sheared Mafic Dyke Predominantly a sheared mafic dyke with up to 30% anorthositic wall-rock inclusions? Overall section carries 15% sulphides as stringers and fracture fillings? Sulphides; 10% Po, 3% Pn and 2% Cpy. Stringers vary from 1mm to 6cm in width and are massive to semi-massive sulphide. Mineralization 85.53 - 87.08 : POPNCP Pyrrhotite/Pentlandite/Chalcopyrite, STR Stringers, 15%							

Hole Number: ES07-56

Units: Metric

Detailed Lithology		Lithology	Assay Data						
From (m)	To (m)		Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
99.60	114.45	<p>ANOR, Anorthosite</p> <p>MIXED ANORTHOSITE - ULTRAMAFIC</p> <p>Predominantly an anorthosite with bands, ribbons and short sections (to 30cm) very dark green ultramafic.</p> <p>Anorthositic component pale cream coloured to pale green and locally buff coloured.</p> <p>Moderately sericitic throughout.</p> <p>Local sections with coarse quartz.</p> <p>Foliation often contorted, but at approximately 80 deg. to CA.</p> <p>Local wispy bright green fuchsite.</p> <p>Minor thin chloritic blebs/bands - dykes?</p> <p>Only longer ultramafic units broken out.</p> <p>Alteration</p> <p>99.60 - 114.45 :CHL Chlorite, H Patchy, W Weak</p> <p>99.60 - 114.45 :Ser Sericite, P Pervasive, M Moderate</p> <p>Structure</p> <p>110.00 - 110.00 : FOL Foliated, 70 Deg to CA</p> <p>MINOR INTERVALS:</p> <p>Minor Interval:</p> <p>104.3 - 104.6 UM, Ultramafic</p> <p>Ultramafic</p> <p>Upper contact irregular - sheared.</p> <p>Very dark green, fine grained and soft</p> <p>Non magnetic.</p> <p>Finely fractured throughout - tectonized.</p> <p>1% to 2% cs. cubic Py.</p> <p>Lower contact at 40 to 45 deg. to CA. - "ragged"</p> <p>Alteration</p> <p>104.30 - 104.60 :SERP Serpentine, P Pervasive, M Moderate</p> <p>Structure</p> <p>104.30 - 104.60 : LC Lower Contact, 40 Deg to CA</p> <p>Minor Interval:</p> <p>104.92 - 105.1 UM, Ultramafic</p> <p>Ultramafic</p> <p>Upper contact at 70 deg. to CA.</p> <p>Similar to previous ultramafic - serpentinous.</p> <p>1% cubic Py</p> <p>Lower contact at 45 to 70 deg. - undulating.</p> <p>Alteration</p> <p>104.92 - 105.10 :SERP Serpentine, P Pervasive, M Moderate</p> <p>Structure</p> <p>104.92 - 104.93 : UC Upper Contact, 70 Deg to CA</p>							

Hole Number: ES07-56

Units: Metric

Detailed Lithology		Assay Data							
From (m)	To (m)	Lithology	Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
		<p>MINOR INTERVALS:</p> <p>Minor Interval: 105.6 - 105.88 UM, Ultramafic Sheared Ultramafic Upper contact at 80 to 85 deg. to CA. Medium to dark green, fine grained and sheared. 1% cubic Py Lower contact at 10 to 35 deg. - "ragged"</p> <p>Alteration 105.60 - 105.88 :SRP Serpentine, P Pervasive, M Moderate</p> <p>Structure 105.60 - 105.61 : UC Upper Contact, 80 Deg to CA</p> <p>Minor Interval: 106.32 - 106.55 UM, Ultramafic Sheared Ultramafic Similar to previous ultramafic - but no Py. Upper contact at 60 deg. to CA - sheared Lower contact at 85 deg. - "ragged".</p> <p>Alteration 106.32 - 106.55 :Sil Silica, P Pervasive, M Moderate</p> <p>Structure 106.32 - 106.33 : UC Upper Contact, 60 Deg to CA 106.55 - 106.55 : LC Lower Contact, 85 Deg to CA</p> <p>Minor Interval: 108.08 - 108.38 MD, Mafic Dike</p> <p>Mafic Dyke Upper contact at 80 deg. to CA Light to medium green and fine grained. Bands/patches brighter green - epidote? Weak to moderate fabric at 65 to 70 deg. to CA. 1% to 2% fine to cs. disseminated Py. Minor fine red-orange mineral - garnet???</p> <p>Lower contact at 80 deg. to CA.</p> <p>Alteration 108.08 - 108.38 :EP Epidote, B Banded, W Weak</p> <p>Structure 108.08 - 108.08 : UC Upper Contact, 80 Deg to CA 108.38 - 108.38 : LC Lower Contact, 80 Deg to CA</p>							

Hole Number: ES07-56

Units: Metric

Detailed Lithology		Lithology	Assay Data						
From (m)	To (m)		Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
114.45	117.50	MD, Mafic Dike SHEARED MAFIC DYKE Upper contact at 70 to 75 deg. to CA - marked by 40mm sheared UM. Mixed medium green - light green - light cream colour and fine grained. Tectonized - mixed with bands/zones anorthositic material. Very minor fine disseminated Py. Lower contact sharp at 70 deg. to CA. Structure 114.45 - 114.45 : UC Upper Contact, 70 Deg to CA 117.50 - 117.50 : LC Lower Contact, 70 Deg to CA							
117.50	124.10	ANOR, Anorthosite SHEARED ANORTHOSITE Predominantly cream coloured with shades of pale green and very pale pink. Coarse bands/patches quartz. Weakly to moderately sericitic throughout. Local weak pervasive hematite alteration Local very dark green bands/ribbons - serpentinous UM inclusions? Local brighter green patches - epidote or related mineral. Unit is not mineralized. Alteration 117.50 - 124.10 :Ser Sericite, P Pervasive, M Moderate 117.50 - 124.10 :HE Hematite, H Patchy, W Weak Structure 120.80 - 120.80 : FOL Foliated, 65 Deg to CA 123.00 - 123.00 : FOL Foliated, 45 Deg to CA							
124.10	126.60	MD, Mafic Dike MAFIC DYKE Upper contact at 70 to 80 deg. to CA. - undulating but sharp. Medium to dark green, fine grained and with a moderately well developed fabric at 75 to 80 deg. to CA. Minor fine white feldspathic bands. Unit is not mineralized. Lower contact sheared at 70 deg. to CA. - 100 grey chill zone? Structure 124.10 - 124.10 : UC Upper Contact, 75 Deg to CA 126.60 - 126.60 : LC Lower Contact, 70 Deg to CA							

Hole Number: ES07-56

Units: Metric

Detailed Lithology		Lithology	Assay Data						
From (m)	To (m)		Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
126.60	143.60	<p>ANOR, Anorthosite</p> <p>SHEARED ANORTHOSITE</p> <p>Very similar to section from 117.50 to 124.10m.</p> <p>Pale cream -green to pink-orange colour.</p> <p>Moderately sericitic throughout.</p> <p>Weak to moderate hematite alteration.</p> <p>Local wispy fuchsitic ribbons.</p> <p>Local very dark green to red-green ultramafic bands/ribbons.</p> <p>Fabric - foliation varies from 40 to 60 deg. to CA.</p> <p>Local very strong buff coloured sericitic bands.</p> <p>Alteration</p> <p>126.60 - 143.60 :Ser Sericite, P Pervasive, M Moderate</p> <p>126.60 - 143.60 :HE Hematite, INT Interstitial, W Weak</p> <p>Structure</p> <p>127.30 - 127.30 : FOL Foliated, 65 Deg to CA</p> <p>132.40 - 132.40 : FOL Foliated, 60 Deg to CA</p> <p>136.10 - 136.10 : FOL Foliated, 70 Deg to CA</p> <p>139.70 - 139.70 : FOL Foliated, 60 Deg to CA</p> <p>141.50 - 141.50 : FOL Foliated, 70 Deg to CA</p> <p>MINOR INTERVALS:</p> <p>Minor Interval:</p> <p>126.9 - 127.2 MD, Mafic Dyke</p> <p>Sheared Mafic Dyke</p> <p>Upper contact at 40 deg. - irregular</p> <p>Medium to dark green and fine grained</p> <p>Well developed fabric at 55 to 60 deg. to CA</p> <p>1% blebby Py - often rimmed by fine hematite.</p> <p>Lower contact undulating at 40 to 60 deg. to CA.</p> <p>Structure</p> <p>126.90 - 127.20 : UC Upper Contact, 40 Deg to CA</p>							
143.60	150.80	<p>MD, Mafic Dyke</p> <p>MAFIC DYKE</p> <p>Upper contact sharp at 75 deg. to CA.</p> <p>Medium to dark green and fine grained.</p> <p>Weak fabric at 75 deg. - marked by alignment of fine feldspars.</p> <p>Several inclusions? anorthositic material.</p> <p>Very minor fine Py and minor streaky and fracture-controlled Po.</p> <p>Lower contact sharp at 75 deg. with a thin grey chill.</p> <p>Structure</p> <p>143.60 - 143.60 : UC Upper Contact, 75 Deg to CA</p> <p>150.80 - 150.80 : LC Lower Contact, 75 Deg to CA</p>							

Hole Number: ES07-56

Units: Metric

Detailed Lithology		Lithology	Assay Data						
From (m)	To (m)		Sample Number	From (m)	To (m)	Length (m)	Ni%	Cu%	Co%
150.80	152.00	ANOR, Anorthosite ANORTHOSITE Mixed pale green to cream coloured and fine grained. Minor sheared mafic component - band/ribbons - dykes? Moderately sericitic throughout. Well developed fabric - foliation - at 70 to 75 deg. to CA. bands/blebs chlorite throughout - alteration? Minor fine speckled biotite Alteration 150.80 - 152.00 :Ser Sericite, P Pervasive, M Moderate 150.80 - 152.00 :CHL Chlorite, H Patchy, W Weak							

Samples

Sample Number	From (m)	To (m)	Ni%	Cu%	Co%
Sample Type	ASSAY				
PG07001	62.55	63.20	0.0050	0.0025	0.0030
PG07002	63.20	63.50	0.0070	0.0060	0.0050
PG07003	63.50	64.62	0.0040	0.0025	0.0030
PG07004	64.62	64.98	1.2250	0.4170	0.0350
PG07005	64.98	65.70	0.0100	0.0025	0.0030
PG07006	84.00	85.00	0.0280	0.0330	0.0030
PG07007	85.00	85.50	0.0120	0.0150	0.0030
PG07008	85.50	85.95	0.7140	0.2390	0.0260
PG07009	85.95	86.40	0.1590	0.1490	0.0100
PG07010	86.40	86.70	1.6400	0.3000	0.0660
PG07012	86.70	87.00	0.2170	0.2780	0.0090
PG07013	87.00	88.00	0.0720	0.1200	0.0050