

Hole Number: ES2004-10

Units: METRIC

|                                  |                                     |                                    |                                |
|----------------------------------|-------------------------------------|------------------------------------|--------------------------------|
| Project Name: Norway - Espedalen | Primary Coordinates Grid: UTM84-32N | Destination Coordinates Grid: UTM: | Collar Dip: -50.00             |
| Project Number: 201              | North: 6803420.18                   | North: 61.36                       | Collar Az: 230.00              |
| Location: Surface                | East: 533158.66                     | East: 9.62                         | Length: 129.65 (m)             |
|                                  | Elev: 1007.65                       | Elev: 1007.65                      | Start Depth: 0.00 (m)          |
| Date Started: Sep 01, 2004       | Collar Survey: Y                    | Plugged: N                         | Contractor: Geo Drilling A/S   |
| Date Completed: Sep 05, 2004     | Multishot Survey: N                 | Hole Size: TT46                    | Core Storage: Strand Fjellstue |
| Logged By: P. Tirschmann         | Pulse EM Survey: Y                  | Casing: Left in Hole, capped       | Final Depth: 129.65 (m)        |

Comments: Purpose: To test UTEM conductor ESP\_16\_16. Conductivity =600 siemens

Result: Intersected variably mineralized, zoned UM body between 50.8m and 83.15m. Conductor can be correlated with the following zones of mineralized ultramafic:

65-73m (8m): 3-15% po disseminations, blebs and stringers in dunite/peridotite

76-79m (3m): 5-10% po blebs, stringers and veinlets in peridotite

Assays: 0.52% Ni, 0.12% Cu, 0.02% Co / 1.00m (65.00-66.00m) (Best)

Borehole UTEM: In-hole response centered @68m.

Lithological interpretation: Anorthositic terrain intruded by mafic dykes and a mineralized ultramafic body.

## Sample Averages

| Average Type | From (m) | To (m) | Length (m) | Ni%    | Cu%    | Co%    |
|--------------|----------|--------|------------|--------|--------|--------|
| WEIGHTED     | 65.00    | 81.00  | 16.00      | 0.3163 | 0.1881 | 0.0188 |

| Detailed Lithology |        | Assay Data  |               |          |        |            |     |     |     |
|--------------------|--------|---|---------------|----------|--------|------------|-----|-----|-----|
| From (m)           | To (m) | Lithology   | Sample Number | From (m) | To (m) | Length (m) | Ni% | Cu% | Co% |
| 0                  | 1.79   | C, Casing   |               |          |        |            |     |     |     |
| 1.79               | 12.60  | MD, Mafic Dike<br>Fine to medium grained, greyish-green mafic dyke. Crosscut by several percent 1-15mm wide white carbonate veinlets. Downhole contact is gradational chilled over approx. 5cm; 60-75 degrees to CA.<br><br>Magnetic Susceptibility: <1<br>Conductivity: Non-conductive<br><br>Interpretation: Unit could be either a gabbroic dyke or a coarse grained mafic metavolcanic flow. (More likely a gabbroic dyke if bounded by anorthosite). Similar units logged as mafic dykes in drillholes ES-2004-05, 06 & 07 (P4 area) and ES-2004-08 & 09 (P2 area).<br>RQD<br>1.79 - 4.00 : 79.00 % RQD 98.00 % Core<br>4.00 - 7.00 : 40.00 % RQD 100.00 % Core<br>7.00 - 10.00 : 68.00 % RQD 100.00 % Core<br>10.00 - 13.00 : 27.00 % RQD 100.00 % Core |               |          |        |            |     |     |     |

Hole Number: ES2004-10

Units: METRIC

| Detailed Lithology |        | Lithology   | Assay Data    |          |        |            |     |     |     |
|--------------------|--------|---|---------------|----------|--------|------------|-----|-----|-----|
| From (m)           | To (m) |   | Sample Number | From (m) | To (m) | Length (m) | Ni% | Cu% | Co% |
| 12.60              | 31.60  | <p>4, Anorthosite / Anorthosite Gabbro</p> <p>White to grey medium grained anorthosite consisting of 80-90% feldspar and 10-20% interstitial soft, green, altered mafic (?) minerals. Local grain size reduction due to shearing. Unit becomes progressively more altered and tectonized downhole. Downhole contact gradational. Cross-cut by 2 narrow ultramafic dykes (see minor intervals).</p> <p>Magnetic Susceptibility: &lt;0.5<br/>Conductivity: Non-conductive</p> <p>Structure<br/>30.40 - 30.41 : Sm General Foliation, 60 Deg to CA<br/>Shearing</p> <p>RQD<br/>13.00 - 16.00 : 72.00 % RQD 100.00 % Core<br/>16.00 - 19.00 : 74.00 % RQD 100.00 % Core<br/>19.00 - 22.00 : 54.00 % RQD 100.00 % Core<br/>22.00 - 25.00 : 64.00 % RQD 100.00 % Core<br/>25.00 - 28.00 : 77.00 % RQD 100.00 % Core<br/>28.00 - 31.00 : 52.00 % RQD 100.00 % Core<br/>31.00 - 34.00 : 48.00 % RQD 95.00 % Core</p> <p>MINOR INTERVALS:<br/>Minor Interval:<br/>23.42 - 24.1 PYXT, Pyroxenite<br/>Fine grained, dark green melano-gabbro to pyroxenite dyke. Contacts sharp; uphole contact at 60 degrees to CA; downhole contact at 50 degrees to CA. Related to uphole gabbroic unit?<br/>Minor Interval:<br/>24.9 - 25.45 PYXT, Pyroxenite<br/>Fine grained, dark green melano-gabbro to pyroxenite dyke. Contacts sharp; uphole contact at 50-55 degrees to CA; downhole contact at approx. 75 degrees to CA. Related to uphole gabbroic unit?</p> |               |          |        |            |     |     |     |

Hole Number: ES2004-10

Units: METRIC

| Detailed Lithology |        | Lithology  | Assay Data    |          |        |            |        |        |        |
|--------------------|--------|--|---------------|----------|--------|------------|--------|--------|--------|
| From (m)           | To (m) |  | Sample Number | From (m) | To (m) | Length (m) | Ni%    | Cu%    | Co%    |
| 31.60              | 50.80  | 4s, Sausseritized/Tectonized Anorthosite   | PG00246       | 49.00    | 50.00  | 1.00       | 0.0250 | 0.0250 | 0.0100 |
|                    |        | Inhomogenous, banded white, grey and pale green sausseritized and sheared anorthosite. Consists of fine grained, sheared sericite and feldspar-rich rock surrounding lozenges of what appears to be coarser grained, equigranular, white anorthosite as uphole. Grain size reduction with shearing evident; locally mylonitized. | PG00247       | 50.00    | 50.80  | 0.80       | 0.0250 | 0.0250 | 0.0100 |
|                    |        | Magnetic Susceptibility: <0.5<br>Conductivity: non-conductive  |               |          |        |            |        |        |        |
|                    |        | Interpretation: Altered, tectonized anorthosite based on gradation to more recognizable anorthosite uphole.  |               |          |        |            |        |        |        |
|                    |        | Structure  |               |          |        |            |        |        |        |
|                    |        | 34.50 - 34.51 : Sm General Foliation, 70 Deg to CA   |               |          |        |            |        |        |        |
|                    |        | 39.70 - 39.71 : Sm General Foliation, 70 Deg to CA   |               |          |        |            |        |        |        |
|                    |        | 43.40 - 43.41 : Sm General Foliation, 70 Deg to CA   |               |          |        |            |        |        |        |
|                    |        | 49.90 - 49.91 : Sm General Foliation, 60 Deg to CA   |               |          |        |            |        |        |        |
|                    |        | RQD  |               |          |        |            |        |        |        |
|                    |        | 34.00 - 37.00 : 64.00 % RQD 100.00 % Core  |               |          |        |            |        |        |        |
|                    |        | 37.00 - 40.00 : 63.00 % RQD 100.00 % Core  |               |          |        |            |        |        |        |
|                    |        | 40.00 - 43.00 : 69.00 % RQD 100.00 % Core  |               |          |        |            |        |        |        |
|                    |        | 43.00 - 46.00 : 67.00 % RQD 100.00 % Core  |               |          |        |            |        |        |        |
|                    |        | 46.00 - 49.00 : 79.00 % RQD 100.00 % Core  |               |          |        |            |        |        |        |
|                    |        | 49.00 - 52.00 : 61.00 % RQD 100.00 % Core  |               |          |        |            |        |        |        |
|                    |        | MINOR INTERVALS:   |               |          |        |            |        |        |        |
|                    |        | Minor Interval:  |               |          |        |            |        |        |        |
|                    |        | 42.65 - 43.03 MD, Mafic Dike   |               |          |        |            |        |        |        |
|                    |        | Pale green, aphanitic mafic dyke. Contacts sharp; uphole contact at 40 degrees to CA; downhole contact at 65 degrees to CA.  |               |          |        |            |        |        |        |

Hole Number: ES2004-10

Units: METRIC

| Detailed Lithology |        | Assay Data   |               |          |        |            |        |        |        |
|--------------------|--------|--|---------------|----------|--------|------------|--------|--------|--------|
| From (m)           | To (m) | Lithology  | Sample Number | From (m) | To (m) | Length (m) | Ni%    | Cu%    | Co%    |
| 50.80              | 83.15  | 6, Undivided Ultramafic Intrusive  | PG00248       | 50.80    | 52.00  | 1.20       | 0.1100 | 0.0700 | 0.0100 |
|                    |        | Fine to medium grained, dark green to black zoned (?) ultramafic body containing trace to, locally, 15% sulphides (po, minor cp) occurring as blebs, disseminations and stringers. Shows less evidence of tectonization compared to surrounding anorthosites. Massive to locally foliated. Several cm to dm scale inclusions (?) of altered, brownish green, fine grained, equigranular gabbroic rock adjacent to downhole contact between 81m and 83.15m. | PG00249       | 52.00    | 53.00  | 1.00       | 0.1200 | 0.0600 | 0.0100 |
|                    |        |  | PG03001       | 53.00    | 54.00  | 1.00       | 0.0600 | 0.0250 | 0.0100 |
|                    |        |  | PG03002       | 54.00    | 55.00  | 1.00       | 0.0800 | 0.0250 | 0.0200 |
|                    |        |  | PG03003       | 55.00    | 56.00  | 1.00       | 0.0900 | 0.0250 | 0.0100 |
|                    |        |  | PG03004       | 56.00    | 57.00  | 1.00       | 0.0700 | 0.0500 | 0.0100 |
|                    |        |  | PG03005       | 57.00    | 58.00  | 1.00       | 0.2300 | 0.0700 | 0.0100 |
|                    |        | 50.8-65m: Fairly soft, pyroxenitic (?) ultramafic largely altered to chlorite, serpentine and talc +- phlogopite. Locally contains mm to cm scale, rounded grains altered to phlogopite (formerly pyroxenes?). Trace disseminated sulphides. Mag sus ranges from <1 to 2.5; non-conductive.  | PG03006       | 58.00    | 59.00  | 1.00       | 0.1400 | 0.0250 | 0.0200 |
|                    |        |  | PG03007       | 59.00    | 60.00  | 1.00       | 0.0250 | 0.0250 | 0.0100 |
|                    |        |  | PG03008       | 60.00    | 61.00  | 1.00       | 0.0250 | 0.0250 | 0.0100 |
|                    |        |  | PG03009       | 61.00    | 62.00  | 1.00       | 0.0250 | 0.0500 | 0.0100 |
|                    |        |  | PG03010       | 62.00    | 62.90  | 0.90       | 0.0250 | 0.0250 | 0.0100 |
|                    |        | 65-73: Medium grained, dark green to black, cumulate textured, well-mineralized dunite/peridotite containing 3-15% sulphide blebs, disseminations & minor fracture fillings (esp. cp). Sulphides locally strung out parallel to foliation. Mag Sus ranges from 2-20; conductivity = 10-30, locally 200 (eg. 65.7m).  | PG03011       | 62.90    | 64.35  | 1.45       | 0.0250 | 0.0250 | 0.0100 |
|                    |        |  | PG03012       | 64.35    | 65.00  | 0.65       | 0.0800 | 0.0250 | 0.0100 |
|                    |        |  | PG03013       | 65.00    | 66.00  | 1.00       | 0.5200 | 0.1200 | 0.0200 |
|                    |        |  | PG03014       | 66.00    | 67.00  | 1.00       | 0.3300 | 0.3100 | 0.0300 |
|                    |        | 73-83.15m: Medium grained, dark green, variably altered peridotitic to pyroxenitic ultramafic containing 1-5% sulphide blebs, disseminations and stringers. Two intervals of fine grained equigranular, melanonorite (?) from 81-81.90m and 82.75-83.15m. Mag sus highly variable (1-50) but averages 15-20; conductivity = 10-60; locally 100-300.  | PG03015       | 67.00    | 68.00  | 1.00       | 0.2600 | 0.1700 | 0.0100 |
|                    |        |  | PG03016       | 68.00    | 69.00  | 1.00       | 0.3000 | 0.2800 | 0.0300 |
|                    |        |  | PG03017       | 69.00    | 70.00  | 1.00       | 0.3700 | 0.3300 | 0.0100 |
|                    |        |  | PG03018       | 70.00    | 71.00  | 1.00       | 0.4200 | 0.3700 | 0.0200 |
|                    |        |  | PG03019       | 71.00    | 72.00  | 1.00       | 0.2900 | 0.1000 | 0.0100 |
|                    |        |  | PG03020       | 72.00    | 73.00  | 1.00       | 0.2000 | 0.0900 | 0.0100 |
|                    |        | Uphole contact sharp at 55 degrees to CA; Downhole contact at approx. 50 degrees to CA.  | PG03021       | 73.00    | 74.00  | 1.00       | 0.1600 | 0.0900 | 0.0300 |
|                    |        |  | PG03022       | 74.00    | 75.00  | 1.00       | 0.2100 | 0.0900 | 0.0200 |
|                    |        |  | PG03023       | 75.00    | 76.00  | 1.00       | 0.2600 | 0.1700 | 0.0100 |
|                    |        | Interpretation: Zoned ultramafic body consisting of a mineralized dunitic to peridotitic core surrounded by peridotitic to pyroxenitic phases. Body interpreted to intrude surrounding anorthosites.   | PG03024       | 76.00    | 77.00  | 1.00       | 0.4700 | 0.3100 | 0.0300 |
|                    |        | Mineralization   | PG03026       | 77.00    | 78.00  | 1.00       | 0.4600 | 0.1100 | 0.0100 |
|                    |        | 50.80 - 53.00 : Po Pyrrhotite, STR Stringers, 0.5%   | PG03027       | 78.00    | 79.00  | 1.00       | 0.2800 | 0.1100 | 0.0200 |
|                    |        | Trace po veinlets locally.   | PG03028       | 79.00    | 80.00  | 1.00       | 0.2300 | 0.1900 | 0.0200 |
|                    |        | 65.00 - 73.00 : Po Pyrrhotite, BB Blebby, 8%   | PG03029       | 80.00    | 81.00  | 1.00       | 0.3000 | 0.1700 | 0.0200 |
|                    |        | 3-15% po, tr cp occurring as disseminations, blebs & stringers   | PG03030       | 81.00    | 81.90  | 0.90       | 0.0900 | 0.1100 | 0.0100 |
|                    |        | 73.00 - 76.00 : Po Pyrrhotite, STR Stringers, 1%   | PG03031       | 81.90    | 82.75  | 0.85       | 0.0800 | 0.0800 | 0.0100 |
|                    |        | 1-2% po, tr cp stringers and blebs   | PG03032       | 82.75    | 83.15  | 0.40       | 0.0600 | 0.0250 | 0.0400 |
|                    |        | 76.00 - 79.00 : Po Pyrrhotite, STR Stringers, 7%   |               |          |        |            |        |        |        |
|                    |        | 5-10% po, tr cp as blebs, stringers and veinlets   |               |          |        |            |        |        |        |
|                    |        | 79.00 - 81.00 : Po Pyrrhotite, BB Blebby, 3%   |               |          |        |            |        |        |        |
|                    |        | 1-5% po, tr py blebs   |               |          |        |            |        |        |        |
|                    |        | 81.00 - 83.15 : Po Pyrrhotite, STR Stringers, 0.5%   |               |          |        |            |        |        |        |
|                    |        | Trace po stringers   |               |          |        |            |        |        |        |
|                    |        | Structure  |               |          |        |            |        |        |        |
|                    |        | 58.30 - 58.31 : Sm General Foliation, 60 Deg to CA   |               |          |        |            |        |        |        |
|                    |        | 67.30 - 67.31 : Sm General Foliation, 45 Deg to CA   |               |          |        |            |        |        |        |
|                    |        | 68.70 - 68.71 : Sm General Foliation, 32 Deg to CA   |               |          |        |            |        |        |        |

Hole Number: ES2004-10

Units: METRIC

| Detailed Lithology |        | Assay Data  |               |          |        |            |        |        |        |
|--------------------|--------|---|---------------|----------|--------|------------|--------|--------|--------|
| From (m)           | To (m) | Lithology   | Sample Number | From (m) | To (m) | Length (m) | Ni%    | Cu%    | Co%    |
|                    |        | RQD<br>52.00 - 55.00 : 69.00 % RQD 100.00 % Core  |               |          |        |            |        |        |        |
|                    |        | 55.00 - 58.00 : 89.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |
|                    |        | 58.00 - 61.00 : 65.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |
|                    |        | 61.00 - 64.00 : 53.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |
|                    |        | 64.00 - 67.00 : 36.00 % RQD 95.00 % Core  |               |          |        |            |        |        |        |
|                    |        | 67.00 - 70.00 : 67.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |
|                    |        | 70.00 - 73.00 : 24.00 % RQD 98.00 % Core  |               |          |        |            |        |        |        |
|                    |        | 73.00 - 76.00 : 45.00 % RQD 98.00 % Core  |               |          |        |            |        |        |        |
|                    |        | 76.00 - 79.00 : 75.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |
|                    |        | 79.00 - 82.00 : 70.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |
|                    |        | 82.00 - 85.00 : 74.00 % RQD 98.00 % Core  |               |          |        |            |        |        |        |
|                    |        | MINOR INTERVALS:<br>Minor Interval:<br>62.9 - 64.35 4s, Sausseritized/Tectonized Anorthosite  |               |          |        |            |        |        |        |
| 83.15              | 110.44 | 4s, Sausseritized/Tectonized Anorthosite  | PG03033       | 83.15    | 84.00  | 0.85       | 0.0250 | 0.0250 | 0.0100 |
|                    |        | Inhomogenous, white, grey and green altered and tectonized anorthosite similar to 31.6-50.8m. Consists of cm to dm scale interval of medium grained, equigranular white and grey anorthosite bounded by much finer grained, foliated, highly sheared anorthosite. Pervasively sausseritized. Rare trace disseminated py (eg. 96m). Appears to be cross-cut by narrow (0.5m) wide serpentized, ultramafic dykes (eg. 89.6-89.9m, 90.1-90.35m, 92.65-93.15m, 94-94.2m, 99-99.35m) between uphole contact and 100m. Ultramafic dykes locally contain trace po. | PG03034       | 84.00    | 85.00  | 1.00       | 0.0250 | 0.0250 | 0.0100 |
|                    |        | Sheared at downhole contact; contact estimated at 80 degrees to CA.   | PG03035       | 87.60    | 88.00  | 0.40       | 0.0700 | 0.0700 | 0.0100 |
|                    |        | Magnetic Susceptibility: <0.50<br>Conductivity: Non-conductive  | PG03036       | 89.00    | 89.50  | 0.50       | 0.0250 | 0.0250 | 0.0100 |
|                    |        | Structure   |               |          |        |            |        |        |        |
|                    |        | 85.70 - 85.71 : Sm General Foliation, 60 Deg to CA  |               |          |        |            |        |        |        |
|                    |        | 98.55 - 98.56 : Sm General Foliation, 60 Deg to CA  |               |          |        |            |        |        |        |
|                    |        | 106.20 - 106.21 : Sm General Foliation, 75 Deg to CA  |               |          |        |            |        |        |        |
|                    |        | RQD   |               |          |        |            |        |        |        |
|                    |        | 85.00 - 88.00 : 62.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |
|                    |        | 88.00 - 91.00 : 82.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |
|                    |        | 91.00 - 94.00 : 84.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |
|                    |        | 94.00 - 97.00 : 54.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |
|                    |        | 97.00 - 100.00 : 76.00 % RQD 100.00 % Core  |               |          |        |            |        |        |        |
|                    |        | 100.00 - 103.00 : 63.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |
|                    |        | 103.00 - 106.00 : 54.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |
|                    |        | 106.00 - 109.00 : 29.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |
|                    |        | 109.00 - 112.00 : 79.00 % RQD 100.00 % Core   |               |          |        |            |        |        |        |

Hole Number: ES2004-10

Units: METRIC

| Detailed Lithology |        | Lithology  | Assay Data    |          |        |            |     |     |     |
|--------------------|--------|--|---------------|----------|--------|------------|-----|-----|-----|
| From (m)           | To (m) |  | Sample Number | From (m) | To (m) | Length (m) | Ni% | Cu% | Co% |
| 110.44             | 129.65 | <p>MD, Mafic Dyke</p> <p>Weakly foliated, homogenous, fine grained, dark green mafic dyke. Crosscut by trace to 2% percent 1-5mm wide white carbonate veinlets. Patchy, pale green bleaching of rock locally (eg. 116.5-120m). Rare trace disseminated pyrite.</p> <p>Magnetic Susceptibility: &lt;1, typically 0.5-1.<br/>Conductivity: Non-conductive</p> <p>Interpretation: Melano-gabbroic dyke.</p> <p>Structure</p> <p>114.20 - 114.21 : Sm General Foliation, 75 Deg to CA<br/>124.20 - 124.21 : Sm General Foliation, 55 Deg to CA<br/>128.65 - 128.66 : Sm General Foliation, 65 Deg to CA</p> <p>RQD</p> <p>112.00 - 115.00 : 96.00 % RQD 100.00 % Core<br/>115.00 - 118.00 : 98.00 % RQD 100.00 % Core<br/>118.00 - 121.00 : 85.00 % RQD 100.00 % Core<br/>121.00 - 124.00 : 97.00 % RQD 100.00 % Core<br/>124.00 - 127.00 : 91.00 % RQD 100.00 % Core<br/>127.00 - 129.65 : 79.00 % RQD 95.00 % Core</p> |               |          |        |            |     |     |     |

## Samples

| Sample Number | From (m) | To (m) | Ni%    | Cu%    | Co%    |
|---------------|----------|--------|--------|--------|--------|
| Sample Type   | ASSAY    |        |        |        |        |
| PG00246       | 49.00    | 50.00  | 0.0250 | 0.0250 | 0.0100 |
| PG00247       | 50.00    | 50.80  | 0.0250 | 0.0250 | 0.0100 |
| PG00248       | 50.80    | 52.00  | 0.1100 | 0.0700 | 0.0100 |
| PG00249       | 52.00    | 53.00  | 0.1200 | 0.0600 | 0.0100 |
| PG03001       | 53.00    | 54.00  | 0.0600 | 0.0250 | 0.0100 |
| PG03002       | 54.00    | 55.00  | 0.0800 | 0.0250 | 0.0200 |
| PG03003       | 55.00    | 56.00  | 0.0900 | 0.0250 | 0.0100 |
| PG03004       | 56.00    | 57.00  | 0.0700 | 0.0500 | 0.0100 |
| PG03005       | 57.00    | 58.00  | 0.2300 | 0.0700 | 0.0100 |
| PG03006       | 58.00    | 59.00  | 0.1400 | 0.0250 | 0.0200 |
| PG03007       | 59.00    | 60.00  | 0.0250 | 0.0250 | 0.0100 |
| PG03008       | 60.00    | 61.00  | 0.0250 | 0.0250 | 0.0100 |
| PG03009       | 61.00    | 62.00  | 0.0250 | 0.0500 | 0.0100 |
| PG03010       | 62.00    | 62.90  | 0.0250 | 0.0250 | 0.0100 |
| PG03011       | 62.90    | 64.35  | 0.0250 | 0.0250 | 0.0100 |
| PG03012       | 64.35    | 65.00  | 0.0800 | 0.0250 | 0.0100 |

Hole Number: ES2004-10

Units: METRIC

## Samples

| Sample Number | From (m) | To (m) | Ni%    | Cu%    | Co%    |
|---------------|----------|--------|--------|--------|--------|
| Sample Type   | ASSAY    |        |        |        |        |
| PG03013       | 65.00    | 66.00  | 0.5200 | 0.1200 | 0.0200 |
| PG03014       | 66.00    | 67.00  | 0.3300 | 0.3100 | 0.0300 |
| PG03015       | 67.00    | 68.00  | 0.2600 | 0.1700 | 0.0100 |
| PG03016       | 68.00    | 69.00  | 0.3000 | 0.2800 | 0.0300 |
| PG03017       | 69.00    | 70.00  | 0.3700 | 0.3300 | 0.0100 |
| PG03018       | 70.00    | 71.00  | 0.4200 | 0.3700 | 0.0200 |
| PG03019       | 71.00    | 72.00  | 0.2900 | 0.1000 | 0.0100 |
| PG03020       | 72.00    | 73.00  | 0.2000 | 0.0900 | 0.0100 |
| PG03021       | 73.00    | 74.00  | 0.1600 | 0.0900 | 0.0300 |
| PG03022       | 74.00    | 75.00  | 0.2100 | 0.0900 | 0.0200 |
| PG03023       | 75.00    | 76.00  | 0.2600 | 0.1700 | 0.0100 |
| PG03024       | 76.00    | 77.00  | 0.4700 | 0.3100 | 0.0300 |
| PG03026       | 77.00    | 78.00  | 0.4600 | 0.1100 | 0.0100 |
| PG03027       | 78.00    | 79.00  | 0.2800 | 0.1100 | 0.0200 |
| PG03028       | 79.00    | 80.00  | 0.2300 | 0.1900 | 0.0200 |
| PG03029       | 80.00    | 81.00  | 0.3000 | 0.1700 | 0.0200 |
| PG03030       | 81.00    | 81.90  | 0.0900 | 0.1100 | 0.0100 |
| PG03031       | 81.90    | 82.75  | 0.0800 | 0.0800 | 0.0100 |
| PG03032       | 82.75    | 83.15  | 0.0600 | 0.0250 | 0.0400 |
| PG03033       | 83.15    | 84.00  | 0.0250 | 0.0250 | 0.0100 |
| PG03034       | 84.00    | 85.00  | 0.0250 | 0.0250 | 0.0100 |
| PG03035       | 87.60    | 88.00  | 0.0700 | 0.0700 | 0.0100 |
| PG03036       | 89.00    | 89.50  | 0.0250 | 0.0250 | 0.0100 |